

q -EULER AND GENOCCHI NUMBERS

TAEKYUN KIM

Institute of Science Education,
Kongju National University, Kongju 314-701, S. Korea
e-mail: tkim@kongju.ac.kr (or tkim64@hanmail.net)

ABSTRACT. Carlitz has introduced an interesting q -analogue of Frobenius-Euler numbers in [4]. He has indicated a corresponding Stadudt-Clausen theorem and also some interesting congruence properties of the q -Euler numbers. In this paper we give another construction of q -Euler numbers, which are different than his q -Euler numbers. By using our q -Euler numbers, we define the q -analogue of Genocchi numbers and investigate the relations between q -Euler numbers and q -analogs of Genocchi numbers.

1. INTRODUCTION

Throughout this paper, we consider a complex number $q \in \mathbb{C}$ with $|q| < 1$ as an indeterminate. The q -analogue of n is defined by $[n]_q = \frac{1-q^n}{1-q}$. The ordinary Euler numbers are defined by the generating function as follows:

$$(1) \quad F(t) = \frac{2}{e^t + 1} = e^{Et} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!}, \quad |t| < \pi,$$

where we use the technique method notation by replacing E^m by E_m ($m \geq 0$), symbolically, cf.[2, 6].

From Eq.(1), we can derive the Genocchi numbers as follows:

$$(2) \quad G(t) = \frac{2t}{e^t + 1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}, \quad |t| < \pi.$$

2000 Mathematics Subject Classification 11S80, 11B68

Key words and phrases: Sums of powers, Bernoulli number, Bernoulli polynomials

It satisfies $G_1 = 1$, $G_3 = G_5 = G_7 = \dots = 0$, and even coefficients are given $G_m = 2(1 - 2^{2m})B_{2m} = 2mE_{2m-1}$, where B_m are the m -th ordinary Bernoulli numbers, cf.[6]. It follows from (2) and Stadudt-Clasusen theorem that Genocchi numbers are integers. For $x \in \mathbb{R}$ (=the field of real numbers) the Euler polynomials are defined by

$$(3) \quad F(x, t) = F(t)e^{xt} = \frac{2}{e^t + 1}e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (|t| < \pi).$$

From (3), we can also derive the definition of Genocchi polynomials as follows:

$$(4) \quad \frac{2t}{e^t + 1}e^{xt} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!}, \quad (|t| < \pi).$$

The following formulae ((5)-(6)) are well known in [6].

$$(5) \quad E_m(x) = \sum_{k=0}^m \binom{m}{k} \frac{G_{k+1}}{k+1} x^{m-k}.$$

For $n, m \geq 1$, and n odd, we have

$$(6) \quad (n^m - n)G_m = \sum_{k=1}^{m-1} \binom{m}{k} n^k G_k Z_{m-k}(n-1),$$

where $Z_m(n) = 1^m - 2^m + 3^m - \dots + (-1)^{n+1}n^m$. In this paper we give the q -analogs of the above Eq.(5) and Eq.(6). The purpose of this paper is to give another construction of q -Euler numbers, which are different than a q -Eulerian numbers of Carlitz. From the definition of our q -Euler numbers, we derive the q -analogs of Genocchi numbers and investigate the properties of q -Genocchi numbers which are related to q -Euler numbers.

2. q -EULER NUMBERS AND POLYNOMIALS

Let q be a complex number with $q < 1$. In [3, 4] Carlitz constructed q -analogue of Eulerian numbers. We now consider another construction of a q -Eulerian numbers, which are different than his q -Eulerian numbers. First we consider the following generating functions:

$$(7) \quad F_q(t) = [2]_q e^{\frac{t}{1-q}} \sum_{j=0}^{\infty} \frac{(-1)^j}{1+q^{j+1}} \left(\frac{1}{1-q} \right)^j \frac{t^j}{j!} = e^{E_q t} = \sum_{n=0}^{\infty} E_{n,q} \frac{t^n}{n!},$$

and

$$(8) \quad F_q(x, t) = [2]_q e^{\frac{t}{1-q}} \sum_{j=0}^{\infty} \frac{(-1)^j q^{jx}}{1+q^{j+1}} \left(\frac{1}{1-q} \right)^j \frac{t^j}{j!} = e^{E_q(x)t} = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!},$$

where we use the technique method notation by replacing E_q^n by $E_{n,q}$, symbolically. Thus we have

$$(8-1) \quad \begin{aligned} E_{n,q} &= \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l}{1+q^{l+1}}, \\ E_{n,q}(x) &= \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} \frac{(-1)^l}{1+q^{l+1}} q^{lx}, \end{aligned}$$

where $\binom{n}{l}$ is binomial coefficient.

By (8-1), we easily see that $\lim_{q \rightarrow 1} E_{n,q} = E_n$ and $\lim_{q \rightarrow 1} E_{n,q}(x) = E_n(x)$. From Eq.(8), we can derive the below Eq.(9):

$$(9) \quad F_q(x, t) = [2]_q \sum_{n=0}^{\infty} (-1)^n q^n e^{[n+x]_q t} = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}.$$

By (9), we easily see that

$$(10) \quad E_{n,q}(x) = \frac{[2]_q}{[2]_{q^m}} [m]_q^n \sum_{a=0}^{m-1} (-1)^a q^a E_{n,q^m} \left(\frac{a+x}{m} \right) = \sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} q^{kx} E_{k,q}, \text{ for } m \text{ odd.}$$

This is equivalent to

$$(11) \quad [2]_{q^m} E_{n,q}(xm) = [2]_q [m]_q^n \sum_{a=0}^{m-1} (-1)^a q^a E_{n,q^m} \left(\frac{a}{m} + x \right), \text{ for } m \text{ odd.}$$

If we put $x = 0$ in Eq.(11), then we have

$$(12) \quad [m]_{-q} E_{n,q} - [m]_q^n \frac{[m(n+1)]_{-q}}{[n+1]_{-q}} E_{n,q^m} = \sum_{l=0}^{n-1} \binom{n}{l} [m]_q^l E_{l,q^m} \sum_{a=1}^{m-1} (-1)^a q^{a(l+1)} [a]_q^{n-l},$$

where $[m]_{-q} = \frac{1+q^m}{1+q}$ for m odd.

Define the operation $*$ on $f_n(q)$ as follows:

$$(13) \quad (1 - [m]_q^n) * f_n(q) = [m]_{-q} f_n(q) - [m]_q^n \frac{[m(n+1)]_{-q}}{[n+1]_{-q}} f_n(q^m).$$

By (12) and (13), we obtain the following:

Proposition 1. *For $m, n \in \mathbb{N}$ and m odd, we have*

$$(1 - [m]_q^n) * E_{n,q} = \sum_{l=0}^{n-1} \binom{n}{l} [m]_q^l E_{l,q^m} \sum_{a=1}^{m-1} (-1)^a q^{a(l+1)} [a]_q^{n-l}.$$

For any positive integer n , it is easy to see that

$$(14) \quad -[2]_q \sum_{l=0}^{\infty} (-1)^{l+n} q^{l+n} e^{[l+n]_q t} + [2]_q \sum_{l=0}^{\infty} (-1)^l q^l e^{[l]_q t} = [2]_q \sum_{l=0}^{n-1} (-1)^l q^l e^{[l]_q t}.$$

From (9) and (14) we can derive the below:

$$\sum_{l=0}^{n-1} (-1)^l q^l [l]_q^m = \frac{1}{[2]_q} ((-1)^{n+1} q^n E_{m,q}(n) - E_{m,q}).$$

Therefore we obtain the following:

Proposition 2. *For $n, m \in \mathbb{N}$, we have*

$$\sum_{l=0}^{n-1} (-1)^l q^l [l]_q^m = \frac{1}{[2]_q} ((-1)^{n+1} q^n E_{m,q}(n) - E_{m,q}).$$

In the recent many authors have studies the sums of powers of consecutive integers, cf.[1, 5, 7, 10, 11]. The above Proposition 2 is the another q -analogue of the sums of powers of consecutive integers. The Genocchi numbers G_n are defined by the generating function:

$$G(t) = \frac{2t}{e^t + 1} = e^{Gt} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}, \quad (|t| < \pi),$$

where we use the technique method notation by replacing G^m by G_m ($m \geq 0$), symbolically. It satisfies $G_1 = 1$, $G_3 = G_5 = G_7 = \dots = 0$ and even coefficients are given $G_m = 2(1 - 2^{2m})B_{2m} = 2mE_{2m-1}$, cf.[6]. We now derive the q -extension of the above Genocchi numbers from the definition of our q -Euler numbers.

3. q -GENOCCHI NUMBERS AND POLYNOMIALS

By the meaning of (1) and (2), let us define the q -extension of Genocchi numbers as follows:

$$G_q(t) = [2]_q t \sum_{n=0}^{\infty} (-1)^n q^n e^{[n]_q t} = \sum_{n=0}^{\infty} G_{n,q} \frac{t^n}{n!}, \quad (|t| < \pi).$$

Note that $\lim_{q \rightarrow 1} G_q(t) = \frac{2t}{e^t + 1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}$. Hence, $\lim_{q \rightarrow 1} G_{n,q} = G_n$. In [8], the q -Bernoulli numbers are defined by

$$(15) \quad -t \sum_{n=0}^{\infty} q^n e^{[n]_q t} = \sum_{n=0}^{\infty} B_{n,q} \frac{t^n}{n!}, \quad (|t| < 2\pi).$$

It was known that $\lim_{q \rightarrow 1} B_{n,q} = B_n$, cf.[8, 9]. By (15), we easily see that

$$(16) \quad -[2]_q t \sum_{n=0}^{\infty} q^n e^{[n]_q t} + 2[2]_q t \sum_{n=0}^{\infty} q^{2n} e^{[2n]_q t} = [2]_q t \sum_{n=0}^{\infty} (-1)^n q^n e^{[n]_q t}.$$

From (15) and (16), we can derive the below Eq.(17):

$$(17) \quad G_{n,q} = [2]_q B_{n,q} - 2[2]_q^n B_{n,q^2}.$$

Let us consider the q -analogue of Genocchi polynomials as follows:

$$(18) \quad G_q(x, t) = [2]_q t \sum_{n=0}^{\infty} (-1)^n q^{n+x} e^{[n+x]_q t} = \sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!}.$$

By (18), we easily see that

$$(19) \quad G_q(x, t) = [2]_q q^{xt} e^{\frac{t}{1-q}} \sum_{l=0}^{\infty} \frac{(-1)^l}{1+q^{l+1}} q^{lx} \left(\frac{1}{1-q} \right)^l \frac{t^l}{l!}.$$

Thus, we have

$$G_{n,q}(x) = n \left(\frac{1}{1-q} \right)^{n-1} \sum_{l=0}^{n-1} \binom{n-1}{l} \frac{(-1)^l}{1+q^{l+1}} q^{(l+1)x}.$$

From (8) and (19), we can derive the below equality:

$$\begin{aligned}
 F_q(x, t) &= [2]_q \sum_{n=0}^{\infty} (-1)^n q^n e^{[n+x]_q t} = \frac{e^{[x]_q t}}{q^x t} [2]_q q^x t \sum_{n=0}^{\infty} (-1)^n q^n e^{q^x [n]_q t} \\
 (20) \quad &= e^{[x]_q t} \sum_{n=0}^{\infty} q^{nx} \frac{G_{n+1,q}}{n+1} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} q^{nx} \frac{G_{n+1,q}}{n+1} \right) \frac{t^n}{n!}.
 \end{aligned}$$

By (20), we easily see that

$$(21) \quad E_{n,q}(x) = \sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} q^{nx} \frac{G_{n+1,q}}{n+1}.$$

Remark. The Eq.(21) is the q -analogue of Eq.(5).

Therefore we obtain the following theorem:

Theorem 3. *For any positive integer n , we have*

$$\begin{aligned}
 (a) \quad G_{n,q}(x) &= n \left(\frac{1}{1-q} \right)^{n-1} \sum_{l=0}^{n-1} \binom{n-1}{l} \frac{(-1)^l}{1+q^{l+1}} q^{(l+1)x}, \\
 (b) \quad E_{n,q}(x) &= \sum_{k=0}^n \binom{n}{k} [x]_q^{n-k} q^{nx} \frac{G_{n+1,q}}{n+1}, \\
 (c) \quad G_{n,q} &= [2]_q B_{n,q} - 2[2]_q^n B_{n,q^2},
 \end{aligned}$$

where $B_{n,q}$ are the q -Bernoulli numbers which are defined in [8].

By (18), we easily see that

$$(22) \quad \sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left(\frac{[2]_q}{[2]_{q^m}} [m]_q^{n-1} \sum_{a=0}^{m-1} (-1)^a q^{a+x} G_{n,q^m} \left(\frac{x+a}{m} \right) \right) \frac{t^n}{n!}, \text{ for } m \text{ odd}.$$

Thus we obtain the following:

Theorem 4. *Let $m \in \mathbb{N}$ and m odd. Then we see that*

$$G_{n,q}(x) = \frac{[2]_q}{[2]_{q^m}} [m]_q^{n-1} \sum_{a=0}^{m-1} (-1)^a q^{a+x} G_{n,q^m} \left(\frac{x+a}{m} \right) = \sum_{k=0}^{\infty} \binom{n}{k} q^{kx} G_{k,q} [x]_q^{n-k}.$$

This is equivalent to

$$(23) \quad G_{n,q}(mx) = \frac{[2]_q}{[2]_{q^m}} [m]_q^{n-1} \sum_{a=0}^{m-1} (-1)^a q^{a+mx} G_{n,q^m}(x + \frac{a}{m}).$$

If we take $x = 0$ in Eq.(23), then we easily see that

$$(24) \quad [2]_{q^m} [m]_q G_{n,q} - [2]_q [m]_q^n G_{n,q^m} \frac{[2]_{q^{m(n+1)}}}{[2]_{q^{n+1}}} = [2]_q \sum_{k=0}^{n-1} \binom{n}{k} [m]_q^k G_{k,q^m} \sum_{a=0}^{m-1} (-1)^a q^{a(k+1)} [a]_q^{n-k}.$$

From the definition of the operation $*$ in the previous section, we note that

$$(25) \quad ([m]_q - [m]_q^n) * f_n(q) = [2]_{q^m} [m]_q f_n(q) - [2]_q [m]_q^n \frac{[2]_{q^{m(n+1)}}}{[2]_{q^{n+1}}} f_n(q^m).$$

By (24) and (25), we easily see that

$$([m]_q - [m]_q^n) * G_{n,q} = [2]_q \sum_{k=0}^{n-1} \binom{n}{k} [m]_q^k G_{k,q^m} \sum_{a=0}^{m-1} (-1)^a q^{a(k+1)} [a]_q^{n-k}.$$

REFERENCES

1. G. E. Andrews, *q-analogs of the binomial coefficient congruences of Babbage, Wolstenhlome and Glaisher*, Discrete Math. **204** (1999), 15-25.
2. T. Apostol, *Introduction to analytic number theory*, Undergraduate Texts in Math., Springer-Verlag, New York, 1986.
3. L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987-1000.
4. L. Carlitz, *q-Bernoulli and Eulerian Numbers*, Trans. Amer. Math. Soc. **76** (1954), 332-350.
5. G. Gasper, M. Rahman, *Basic hypergeometric series*, Encyclopedia of Mathematics and Its Applications, Vol 96, Secod Edition, Cambridge Univ. Press, 2004.
6. F. T. Howard, *Applications of a recurrence for the Bernoulli numbers*, J. Number Theory **52** (1995), 157-172.
7. T. Kim, *Sums of powers of consecutive q-integers*, Advan. Stud. Contemp. Math. **9** (2004), 15-18.
8. T. Kim, *Analytic continuation of multiple q-zeta functions and their values at negative integers*, Russian J. Math. Phys. **11** (2004), 71-76.
9. T. Kim, *Multiple p-adic L-Function*, Russian J. Math. Phys. **13** no. 1 (2006), 00-00.
10. D. E. Knuth, *Johann Faulhaber and sums of powers*, Math. Comput. **61** (1993), 277-294.
11. Y.-Y. Shen, *A note on the sums of powers of consecutive integers*, Tunghai Science **5** (2003), 101-106.